CHECKING IMMUNIZATION RECORDS

- 1. Determine the age of the child at the time of inspection.
- 2. Compare the child's immunization record with the chart.
- 3. Determine if the correct number of doses for the required vaccinations have been administered.
 - A. If yes, they are in compliance.
 - B. *If no*, ask the provider if the child is conditionally enrolled.
 - i. If yes, check for the first dose of required vaccinations, and if the child is on schedule. The child is in compliance if they have received the first round of doses and are current with their catch-up schedule. The child is out of compliance if they have not received the first dose, or if they are more than 30 days behind the catch-up schedule.
 - ii. *If no*, it is a finding.

Number of Doses Required for Age of the Child

	3-4 months	5-6 months	7-12 months	13-15 months	16-18 months	19-59 months
DTaP / DTP (Diphtheria, Tetanus, Pertussis)						
Brand Names: ACEL-IMUNE, ADACEL, Boostrix, Certiva, DT or Td (diptheria & tetanus only), Infanrix, PEDIARIX, Pentacel, TETRAMUNE, TriHIBIT, Tripedia	1	2	3	3	3	4
* Hib (Haemophilus Influenzae type b)						
Brand Names: ActHIB, HibTITER, OmniHIB, Pentacel, ProHIBIT, TETRAMUNE, TriHIBIT,	1	2	3**	3**	4**	4**
COMVAX, PedvaxHIB						
Polio	1	2	2	2	2	3
Brand Names: IPV, IPOL, OPV, ORIMUNE, PEDIARIX, Pentacel, Sabin, Salk	·	_	_	_	_	Ů
MMR (Mumps, Measles, Rubella)						
Brand Names: Attenuvax (measles only), Meruvax II (rubella only), MR (measles & rubella), M-M-R-	0	0	0	0	1	1
II, Mumpsvax (mumps only), MMRV, Proquad						
Hep B (Hepatitis B)	1	2	2	2	2	3
Brand Names: COMVAX, Engerix-B, PEDIARIX, RECOMBIVAX HB	'			_		Ŭ
Varicella (Chickenpox)	0	0	0	0	0	1
Brand Names: VARIVAX, MMRV, Proquad	U	U	U	U	U	l
Hep A (Hepatitis A)	0	0	0	1	1	1
Brand Names: Havrix, VAQTA	U	U	U		I	l
* PCV (Pneumococcal)	1	2*	2*	3*	1*	1*
Brand Names: Prevnar	I		3*	3	4*	4*

^{*} May be in compliance with 1-3 doses of this vaccine. See delayed schedule charts to determine if the correct number of doses have been received.

^{**} In compliance with one less dose, If first two doses received were COMVAX and/or PedvaxHIB.